



## Post-remagnetisation vertical axis rotation and tilting of the Murihiku Terrane (North Island, New Zealand)

Suzanna H. A. van de Lagemaat, Lydian M. Boschman, Peter J. J. Kamp, Cor G. Langereis & Douwe J. J. van Hinsbergen

To cite this article: Suzanna H. A. van de Lagemaat, Lydian M. Boschman, Peter J. J. Kamp, Cor G. Langereis & Douwe J. J. van Hinsbergen (2017): Post-remagnetisation vertical axis rotation and tilting of the Murihiku Terrane (North Island, New Zealand), *New Zealand Journal of Geology and Geophysics*, DOI: [10.1080/00288306.2017.1400983](https://doi.org/10.1080/00288306.2017.1400983)

To link to this article: <https://doi.org/10.1080/00288306.2017.1400983>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



[View supplementary material](#)



Published online: 30 Nov 2017.



[Submit your article to this journal](#)



Article views: 11



[View related articles](#)



[View Crossmark data](#)

## Post-remagnetisation vertical axis rotation and tilting of the Murihiku Terrane (North Island, New Zealand)

Suzanna H. A. van de Lagemaat<sup>a</sup>, Lydian M. Boschman<sup>a</sup>, Peter J. J. Kamp<sup>b</sup>, Cor G. Langereis<sup>a</sup> and Douwe J. J. van Hinsbergen<sup>a</sup>

<sup>a</sup>Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands; <sup>b</sup>School of Science, University of Waikato, Hamilton, New Zealand

### ABSTRACT

We collected palaeomagnetic sample sets from Murihiku Terrane, North Island to constrain its palaeolatitude during the Late Triassic–Jurassic. The majority of the sample host rocks were remagnetised. However, a few samples show a magnetic signal that possibly recorded a primary remanent magnetisation. These samples preliminarily indicate that Murihiku Terrane was located at c. 63°S during the Early Jurassic. The remagnetised samples reveal significant post-remagnetisation tectonic rotation and tilting of the host rocks. We estimated an  $83 \pm 5$  Ma timing of remagnetisation by plotting the palaeolatitude data on the apparent polar wander path of northern Zealandia. Samples from southernmost sites have lower inclination, which we interpret as reflecting eastward post-remagnetisation tilt of this region by 20°. In addition, declination data indicate large-scale post-remagnetisation rotation of Port Waikato and Awakino Gorge areas. This study contributes to the ongoing debate on the age and tectonic origin of oroclines in New Zealand basement.

### ARTICLE HISTORY

Received 12 April 2017  
Accepted 2 November 2017

### ASSOCIATE EDITOR

Associate Professor Andrew Gorman

### KEYWORDS

Paleomagnetism; New Zealand; Murihiku Terrane; Kawhia syncline; Gondwana

### Introduction

The New Zealand continent (Zealandia) separated from Gondwana during the Late Cretaceous following continental rifting and sea floor spreading in Tasman Sea and between Campbell Plateau and Marie Byrd Land (MacKinnon 1983; Haston and Luyendyk 1991; Mortimer 2004). The majority of basement on land in New Zealand, the so-called Eastern Province, consists of volcanic arc (Brook Street Terrane), ocean floor (Dun Mountain Terrane), forearc (Murihiku Terrane) and accretionary wedge elements (Torlesse complex), interpreted to have formed at one or more intra-oceanic subduction zones (Mortimer et al. 2014). This allochthonous assemblage is separated from Gondwana-derived units in western New Zealand (North-west Nelson, Westland, Fiordland) by the Median Batholith (c. 375 and 110 Ma), which represents a long-lived magmatic arc along the Gondwana margin (Mortimer 2004).

The Murihiku Terrane, noted above is one of the Eastern Province elements, is structurally separate from adjacent terranes, forming a simple yet long syncline, known in the North Island as the Kawhia syncline and in the South Island as the Southland syncline. This terrane has long been regarded as a Triassic–Jurassic forearc basin based on its simple

structure, its well-bedded and fossil-rich character and its volcanic arc provenance (Briggs et al. 2004).

Here, we present the results of a paleomagnetic study of rocks within the Kawhia syncline in an attempt to constrain the paleolatitude of the Murihiku Terrane during the Jurassic, and hence to assess the amount of probable Early Cretaceous convergence between it and Paleozoic terranes comprising eastern Gondwana. We are aware that previous paleomagnetic studies have concluded widespread remagnetisation of the Eastern Province basement of New Zealand (Haston and Luyendyk 1991), including within the Kawhia syncline (Oliver 1994; Kodama et al. 2007). Nevertheless, we hypothesise that the analysis of samples from different facies and localities within the Murihiku Terrane may identify some that retain a detrital remanent magnetisation from which the paleolatitude can be calculated. Even if the rocks drilled at all sampled sites turn out to be remagnetised, these paleomagnetic data may be useful to help constrain the post-remagnetisation tectonic history of the terrane.

### Geological setting

The Murihiku Terrane mostly comprises Triassic to Upper Jurassic marine and non-marine sediments

**CONTACT** Suzanna H.A. van de Lagemaat  [suzannavdl@gmail.com](mailto:suzannavdl@gmail.com)

 Supplemental data for this article can be accessed <https://doi.org/10.1080/00288306.2017.1400983>.

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

(Ballance and Campbell 1993) consisting mainly of volcanoclastic sandstone and siltstone with conglomerate in places and frequent tuff layers (Aita and Spörli 1992). These strata span c. 90 Ma and have thicknesses of up to 10 km or more (Coombs et al. 1992; Roser et al. 2002). Zeolite facies have formed through low-grade burial metamorphism (Coombs 1954; Ballance et al. 1980; Roser et al. 2002).

Petrographic and geochemical studies of Murihiku Terrane samples show that they derived from direct volcanic eruptive materials and mainly from erosion of a volcanic arc terrane (Roser et al. 2002; Briggs et al. 2004). The sediments and tuff accumulated within a forearc basin probably underlain by oceanic crust (Spörli 1987; Aita and Spörli 1992). However, an origin as an intra-arc or back-arc basin has also been suggested (Howell 1980; Coombs et al. 1992). A paleomagnetically determined paleolatitude for Murihiku basin strata would help constrain the intra-Panthalassa Ocean location where the volcanic arc may have formed.

A previous paleomagnetic study of Lower Jurassic Murihiku Terrane volcanic rocks in the Southland syncline suggested that the basin was located at a high paleolatitude, close to the eastern Gondwana margin (Grindley et al. 1980). The similarity between Triassic plant fossils in Murihiku Terrane strata and from nearby coastal regions of Gondwana has also been used to suggest that the origin of the Murihiku basin must have been close to the Gondwana margin (Retallack 1987). However, the presence of two synchronous magmatic/volcanic arcs of the Median Batholith and the Brook Street Terrane in particular, has been used to suggest that a subduction plate boundary must have existed between the intra-oceanic arc and Gondwana during the Permian prior to accumulation of sediments now exposed in the Murihiku Terrane (Bradshaw 1994; Mortimer 2004). Whether the Murihiku Terrane basin originated adjacent to or at a distance from the eastern Gondwana margin thus remains uncertain, but all tectonostratigraphic terranes were sutured along that margin by the end of the Early Cretaceous (Howell 1980).

At present, the Murihiku Terrane is exposed in both North and South Islands as a broad synclinorium. From south to north, the synclinorium consists of the Southland, Key Summit and Nelson synclines in the South Island and the Kawhia syncline in the North Island (Campbell et al. 2003). The Murihiku Terrane synclinorium is dextrally offset by c. 460 km along the Alpine Fault (e.g. Edbrooke et al. 2015). Regional strikes of Murihiku and other basement bends towards the Alpine Fault and form large oroclines on either side of this fault. The age of oroclinal bending within these terranes as well as its relation to Alpine Fault deformation remains uncertain. A widely held view is that Cenozoic plate motion within New Zealand was accommodated across the Alpine Fault as well as in a broad zone of deformation, leading to

bending of basement terranes as a result of large-scale drag (Kamp 1987; Sutherland 1999; Mortimer 2014). An alternative view is that the basement terranes were bent pre-Cenozoic and that the curvature of the terranes remained unchanged in the Cenozoic because Neogene dextral displacement on Alpine Fault occurred in a very narrow zone (Lamb et al. 2016).


In the Kawhia syncline, the location of our paleomagnetic sampling, the Murihiku Terrane is exposed along the west coast of the central-western North Island between Port Waikato in the north and Awakino Gorge in the south as well as inland. The Kawhia syncline is c. 40 km wide and the strike of its fold axis varies from north-northeast in the south to north-northwest in the north (Figure 1). The age of Murihiku Terrane strata within the Kawhia syncline varies from Late Triassic (Norian) to Late Jurassic (Tithonian) (Roser et al. 2002).

## Methods

Rock samples from Murihiku Terrane were obtained from three locations (Figure 1, Table 1) using a petrol-powered drill with a drill bit having an internal diameter of 25 mm. We always sampled over a sufficiently long interval (10–20 m) per site, enough to average out paleosecular variation (see A95 parameter in Table 2 and explanation below). The cores were oriented in the field using a Brunton magnetic compass with an inclinometer attached. Cores were cut into subsamples 22 mm long using a double-blade circular saw.

Laboratory analyses were carried out at the paleomagnetic laboratory, Fort Hoofddijk, at Utrecht University, the Netherlands. The natural remanent magnetisation of samples was measured on a 2G DC-SQUID magnetometer and further investigated using thermal as well as alternating field (AF) stepwise demagnetisation. Thermal demagnetisation was carried out using successively higher temperatures (in °C): 20, 100, 150, 180, 210, 240, 270, 300, 330, 360, 390, 420, 450, 480, 500, 520, 540, 560 and 580. During AF demagnetisation we used the following field strengths (in mT): 0, 4, 8, 12, 16, 20, 25, 30, 35, 40, 45, 50, 60, 70, (80, 90, 100). Thermomagnetic analyses to determine the nature of magnetic carriers were performed on representative samples for each locality using a horizontal translation-type Curie balance with cycling applied magnetic field, usually 150–300 mT (Mullender et al. 1993). We used (finely crushed) bulk material, usually 50–80 mg. A number of heating-cooling cycles was applied to detect magneto-mineralogical alterations during heating. We used the following temperature scheme (in °C): 20–150, 100–250, 150–350, 250–400, 350–500 and 400–700.

Statistical analysis and interpretation were performed using the online, platform independent portal: [www.paleomagnetism.org](http://www.paleomagnetism.org) (Koymans et al. 2016). Demagnetisation diagrams are plotted as orthogonal vector diagrams (Zijderveld 1967). Interpretation of



**Figure 1.** Simplified geological map of Murihiku Terrane exposed within Kawhia syncline (based on Edbrooke 2001, 2005). Note the change in strike of the fold axis (dashed line) of the syncline from north–northeast in the south to north–northwest in the north. Stars indicate sample localities. Inset, Map of North Island showing the distribution of Murihiku Terrane (orange colour) (based on Mortimer 2004).

demagnetisation diagrams was performed by determining a characteristic remanent magnetisation (ChRM) for components decaying towards the origin. We determined great circles if we found no clear ChRM decaying towards the origin because of a pervasive (low temperature or low coercive) overprint causing overlapping blocking temperatures or coercivity. Lines (ChRM, denoted as ‘setpoints’) and planes (great circles) were determined following an eigenvector approach (Kirschvink 1980). If we have both setpoints and great circles in

a site, we use the method of McFadden and McElhinny (1988) to determine great circle solutions. We applied a  $45^\circ$  cut-off to the virtual geomagnetic pole (VGP) distribution of a set of directions (following Johnson et al. 2008; Deenen et al. 2011). This is an arbitrary fixed angle cut-off meant to remove outliers due to excursions or transitional directions, or to remove outliers due to (assumed, possible) errors in sampling and orientation measurement. Mean directions were determined using Fisher (1953), whereas directional statistics were derived from the corresponding VGP distribution (Deenen et al. 2011), and errors in declination ( $\Delta D_x$ ) and inclination ( $\Delta I_x$ ) were calculated from the cone of confidence (A95) of the mean VGP following Butler (1992). We applied the reliability criteria of Deenen et al. (2011) by determining A95 of the VGP distribution, and calculate the N-dependent values of  $A95_{\min}$  and  $A95_{\max}$  (recalculated by Deenen et al. 2014). Values plotting within this envelope can be explained by paleosecular variation. Values of  $A95 > A95_{\max}$  may indicate additional sources of scatter, while values of  $A95 < A95_{\min}$  represent low dispersion (high  $k$ -values, as with lavas) and cannot reliably represent paleosecular variation. To test the primary origin of the ChRM and if the data permit, we perform field tests. The fold test follows an eigenvector approach (Tauxe and Watson 1994) or the reversal test using the coordinate bootstrap test (Tauxe et al. 2010). The results are compared with the expected directions for a reference locality from the Global Apparent Polar Wander Path of Torsvik et al. (2012). All methods used (and more) are available at [www.paleomagnetism.org](http://www.paleomagnetism.org). In the online supplementary information, we provide all the demagnetisation results and interpretations as a *.dir* file that can be easily imported into [www.paleomagnetism.org](http://www.paleomagnetism.org). Similarly, all statistical results are provided as a *.pmag* file.

## Location descriptions

### Awakino Gorge

The southernmost area we sampled was a road section exposed in Awakino Gorge (AG), (Figure 1, Table 1).

**Table 1.** Information about GPS location of the site, measured bedding plane (strike/dip) and their age plus reference and the number of demagnetised samples per site.

Site	Strike/Dip	GPS coordinates		Age (stage)	Age (Ma)	Reference	No. of samples demagnetised	
		Latitude (N)	Longitude (E)				Th.	AF
AG1	302/63	−38.6739	174.6876	Middle Rhaetian	207–204	Hay (1967)	5	12
AG2	026/70	−38.6737	174.7108	Middle Toarcian	181–176	Hay (1967)	7	22
AG3	022/54	−38.6270	174.7309	Bajocian	170–168	Hay (1967)	5	28
AG4	021/60	−38.6370	174.7275	Bajocian	170–168	Hay (1967)	3	10
KI1	327/33	−38.3268	174.7041	Lower Rhaetian	209–207	Kear (1960)	8	26
KI2	328/37	−38.3168	174.7108	Upper Rhaetian	204–201	Kear (1960)	5	17
KI3	310/29	−38.3201	174.7075	Middle Rhaetian	207–204	Kear (1960)	3	14
KI4	336/29	−38.3201	174.7075	Middle Rhaetian	207–204	Kear (1960)	3	13
KI5	323/29	−38.3201	174.7075	Middle Rhaetian	207–204	Kear (1960)	6	26
PW1	175/36	−37.3993	174.7108	Late Tithonian	147–145	Challinor (2001)	8	40

**Table 2.** Geomagnetic directions per site.

Site	Ns	Na	D	$\Delta D_x$	I	$\Delta I_x$	K	A95 <sub>min</sub>	A95	A95 <sub>max</sub>	$\lambda$
Geographic coordinates											
AG1	17	16	130.3	12.3	-63.8	6.8	19.5	4.0	8.6	14.3	-45.5
AG2	29	21	103.4	5.6	-49.7	5.4	44.6	3.6	4.8	12.0	-30.5
AG3	34	31	98.3	8.4	-62.4	5.0	19.2	3.0	6.0	9.4	-43.8
AG4	13	13	81.0	9.0	-55.0	7.2	33.4	4.3	7.3	16.3	-35.5
AG all	93	79	101.9	5.0	-58.5	3.5	17.7	2.1	3.9	5.2	-39.2
<b>AG E20</b>	<b>85</b>	<b>77</b>	<b>121.0</b>	<b>12.0</b>	<b>-77.0</b>	<b>2.9</b>	<b>11.4</b>	<b>2.1</b>	<b>5.0</b>	<b>5.3</b>	<b>-65.3</b>
AG2*	8	8	301.3	5.4	5.8	10.7	106.6	5.2	5.4	22.1	2.9
AG2* E20	8	8	304.1	6.2	22.7	10.9	83.3	5.2	6.1	22.1	11.8
KI1	33	30	37.3	18.1	-78.6	3.7	16.6	3.1	6.7	9.6	-68.0
KI2	22	20	60.5	11.3	-70.0	4.5	25.4	3.6	6.6	12.4	-54.0
KI3	17	17	356.2	12.5	-74.0	3.8	34.1	3.9	6.2	13.8	-60.1
KI4	16	15	17.3	10.7	-71.7	3.8	43.6	4.1	5.9	14.9	-56.5
KI5	32	30	14.9	8.1	-69.9	3.3	30.6	3.1	4.8	9.6	-53.1
<b>KI all</b>	<b>121</b>	<b>116</b>	<b>26.8</b>	<b>6.9</b>	<b>-74.4</b>	<b>2.0</b>	<b>16.3</b>	<b>1.8</b>	<b>3.4</b>	<b>4.1</b>	<b>-60.7</b>
<b>PW1</b>	<b>48</b>	<b>44</b>	<b>346.2</b>	<b>9.5</b>	<b>-74.1</b>	<b>2.9</b>	<b>22.2</b>	<b>2.6</b>	<b>4.7</b>	<b>7.6</b>	<b>-60.3</b>
Tectonic coordinates											
AG1	17	17	184.9	8.1	-36.4	13.4	21.5	3.9	7.9	13.8	-13.9
AG2	29	21	311.8	7.9	-58.7	5.5	28.0	3.6	6.4	12.0	-39.4
AG3	34	32	305.9	8.0	-61.2	5.0	19.6	3.0	5.9	9.2	-42.3
AG4	13	13	322.9	10.7	-57.2	7.9	25.2	4.3	8.4	16.3	-37.8
AG all	93	65	310.5	4.9	-59.7	3.3	23.7	2.3	3.7	5.9	-40.5
AG E20	N/A										
AG2*	8	8	316.8	27.0	74.9	7.5	20.6	5.2	12.5	22.1	61.6
<b>AG2* E20</b>	<b>8</b>	<b>8</b>	<b>308.3</b>	<b>27.6</b>	<b>75.2</b>	<b>7.5</b>	<b>20.6</b>	<b>5.2</b>	<b>12.5</b>	<b>22.1</b>	<b>62.1</b>
KI1	33	33	247.0	10.8	-67.1	5.1	13.9	3.0	7.0	9.1	-49.9
KI2	22	22	237.9	13.0	-70.4	5.0	18.1	3.5	7.5	11.7	-54.6
KI3	17	17	253.3	8.5	-69.6	3.5	50.2	3.9	5.1	13.8	-53.4
KI4	16	15	286.3	8.4	-63.6	3.6	55.4	4.1	5.2	14.9	-51.8
KI5	32	29	276.3	8.4	-71.7	3.0	34.7	3.1	4.6	9.8	-56.5
KI all	121	119	257.3	5.5	-70.3	2.1	17.6	1.8	3.2	4.0	-54.3
PW1	48	48	61.0	4.9	-54.6	4.0	27.0	2.6	4.0	7.2	-35.1

Ns=total number of measured samples, Na=number of samples after 45° cut-off

Declination/inclination (D, I), their respective errors ( $\Delta D_x$ ,  $\Delta I_x$ ) and paleolatitude ( $\lambda$ ) in degrees

AG2 includes samples taken at AG2, excluding AG2\*, which is the sample set that has a deviating magnetic signal and is listed separately. Values given in bold are used for our paleomagnetic analysis and interpretation. AG E20 refers to the Awakino Gorge sample set corrected for the 20° eastward dip.

The oldest rocks (Site AG1) we drilled are of Middle Rhaetian age (207–204 Ma, Otapiian New Zealand Stage) followed upwards by Middle Toarcian Site AG2 (181–176 Ma, upper Ururoan to lower Te Temaikian New Zealand Stage) and then Bajocian Sites AG3 and AG4 (170–168 Ma, mid-Temaikian New Zealand Stage) (Hay 1967). Stage boundary ages are based on the geological time scale of Gradstein et al. (2012). The 94 samples collected include both sandstone and siltstone (mudstone) facies. Cores from AG1 are mostly quite weathered. At AG2, some cores comprise very fine bluish-grey siltstone and some comprise sandstone. Cores from AG3 comprise bluish-grey siltstone of which the top 0.5 cm is weathered to a brownish colour. The cores taken from AG4 are from two turbidite units comprising silty sandstone.

### Kiritehere

Some 87 cores were collected from five sites at Kiritehere Beach (KI) (Figure 1, Table 1), these sites being accessible only during low tide. One drill site (KI1) is located on the shore platform south of Rararimu Stream, and the four other sites (KI2–5) lie on the shore platform north of this stream. The cores comprise bluish-grey siltstone that was easy to drill. The age of the cores obtained from Kiritehere range from lowermost Rhaetian (209–207 Ma, Otapiian New Zealand stage, Site

KI1) to Middle Rhaetian (207–204 Ma, Site KI3–5) and Upper Rhaetian (204–201 Ma, Site KI2) (Kear 1960). After cutting, 116 specimens were available for paleomagnetic analysis.

### Port Waikato

The most northerly area sampled within the Kawhia syncline was along the rocky coast immediately south of Port Waikato (PW), which is mainly accessible during low tide. This section comprises central parts of the Kawhia syncline and some of the youngest beds preserved within it (Late Tithonian, c. 147–145 Ma; Puarooan New Zealand Stage) (Challinor 2001) (Figure 1, Table 1). At Port Waikato, the sedimentary succession comprises Coleman Conglomerate (a formational unit), which consists of siltstone, sandstone and minor conglomerate (Challinor 2001). The 24 cores we drilled comprise mainly silty sandstone, yielding 48 specimens for paleomagnetic analysis.


## Results

### Magnetic carriers


We carried out thermomagnetic analyses on selected samples at each locality and present representative examples in Figure 3. From Awakino Gorge, sample

AG1.2 (Figure 2A) shows magnetisation that has an almost reversible decrease until 580 °C (Figure 2A), indicating magnetite as the main magnetic carrier. At temperatures above 350 °C, however, there is a slight decrease in the cooling curves, which suggests some contribution of maghemite that inverts to hematite at c. 350 °C (Dankers 1978). The final cooling curve is lower on the figure indicating that at least part of the magnetite/maghemite has been oxidised. Sample AG3.5 (Figure 2B) shows similar behaviour, but above 420 °C there is a small increase followed by a higher cooling curve. This indicates the new formation of magnetite from (a small amount of) pyrite that inverts to magnetite at temperatures above 390–420 °C, which subsequently oxidises to maghemite and finally to haematite, causing the increase and

subsequent decrease in magnetisation (Passier et al. 2001). This is visible in thermomagnetic curves. In thermal demagnetization, the newly formed magnetite may produce spurious demagnetisation behaviour at higher temperatures. Samples from Kiritehere show a significant increase in magnetisation above c. 420 °C, which is caused by the transformation of pyrite into magnetite. The newly formed magnetite is subsequently demagnetised at temperatures above 500 °C. In sample KI1.21 (Figure 2C) there is again a slight loss of magnetisation in the cooling curves after heating to 350 °C, indicating the presence of maghemite. The final cooling curve is lower, meaning that the available maghemite and magnetite have been oxidised. In sample KI2.10 (Figure 2D), the newly formed magnetite from pyrite is (nearly) fully demagnetised. There



**Figure 2.** Representative thermomagnetic curves from a Curie balance using a cycling field between 50 and 300 mT (Mullender et al. 1993). Heating is applied to a maximum temperature of 700 °C in a number of heating–cooling cycles (thin red lines) with the final cooling curve as a thick blue line. Samples are from Awakino Gorge (AG), Kiritehere (KI) and Port Waikato (PW). We used crushed bulk material (typically 50–80 mg). Interpretations are described in the text.



**Figure 3.** Representative examples of demagnetisation diagrams (Zijderveld 1967) in geographic coordinates from our three sample localities (AG, KI and PW).  $I_{NRM}$  indicates the initial natural remanent magnetisation, closed (open) symbols are projections on the horizontal (vertical) plane, and the respective projections (green, red lines) denote the interpreted characteristic remanent magnetisation (ChRM). Numbers refer to temperature ( $^{\circ}\text{C}$ ) or AF field (mT) steps. The initial natural remanent magnetisation intensities ( $I_{NRM}$ ) are given in mA/m.

is no notable contribution of maghemite considering the final cooling curve, which is only marginally higher than the heating curves, meaning that there are some traces left of the newly formed magnetite.

Samples from Port Waikato generally show behaviour similar to those of Kiritehere (PW1.24B, Figure 2E). In sample PW1.2 (Figure 2F) there is a different behaviour; above 450  $^{\circ}\text{C}$  some new magnetite is

formed but it is again demagnetised at c. 500 °C. Yet the final cooling curve is significantly higher than the heating curves, and new magnetic material is probably formed at temperatures up to 700 °C. This is possibly magnetite, judged from the breakdown of clay minerals at the highest temperature reached (700 °C).

### Demagnetisation behaviour and paleomagnetic directions

Samples from Awakino Gorge that were thermally demagnetised generally show the removal of a low-temperature or low-coercive component, followed by a remanent magnetisation component that is stable and decays linearly to the origin above 180–210 °C or above 12 mT. This component is completely removed in most samples at 500 °C (Figure 3A), or at c. 60 mT (Figure 3B), which we interpret as the ChRM.

Mean paleomagnetic directions for all sites are listed in Table 2. ChRMs interpreted from Awakino Gorge samples show variation in mean declination from 81.0° (AG4) to 130.3° (AG1) (Table 2). In geographic coordinates, mean inclinations vary between –49.7° (AG2) and –63.8° (AG1) (Table 2). When all AG sites are grouped together, 14 of 93 directions are rejected by the 45° cut-off (Figure 4A,B). All samples indicate normal polarity and give a mean direction with  $D = 101.9 \pm 5.0^\circ$  and  $I = -58.5 \pm 3.5^\circ$  (where  $D$  = declination,  $I$  = inclination). After tilt correction, the mean direction is  $D = 310.5 \pm 4.9^\circ$ ;  $I = -59.7 \pm 3.3^\circ$  (Figure 4A,B, Table 2).

Remarkably, eight samples form a separate cluster with very low inclinations before bedding tilt correction (see yellow symbols, Figure 4). These samples were all taken at Toarcian site AG2. These samples do not show characteristic differences in composition compared to the other samples taken at AG2. Two of these samples were thermally demagnetised and both have a stable natural remanent magnetisation component between 300 and 500 °C. The stable natural remanent magnetisation component during AF demagnetisation is obtained between 30 and 60 mT (Figure 5). The mean *in situ* direction of these samples (AG2\*) is  $D = 301.3 \pm 5.4^\circ$  and  $I = 5.8 \pm 10.7^\circ$ , and a tilt-corrected direction  $D = 316.8 \pm 27.0^\circ$  and  $I = 74.9 \pm 7.5^\circ$  (Table 2).

Thermal demagnetisation behaviour in Kiritehere samples generally shows a stable natural magnetisation component between 300 and 500 °C, and occasionally between 180 and 500 °C (Figure 3C). AF demagnetisation diagrams show linear trends and the magnetisation component is generally removed at 60–80 mT (Figure 3D).

Because of the relatively small stratigraphic range from which the sites (KI1–5) were taken at Kiritehere, directions from all 116 measured specimens were

grouped and treated as a single set for statistical analysis (Figure 4C,D, Table 2). Only five samples are rejected by the 45° cut-off. All sites indicate normal polarity and give a direction  $D = 26.8 \pm 6.9^\circ$  and  $I = -74.4 \pm 2.0^\circ$ , and a tilt-corrected direction  $D = 257.3 \pm 5.5^\circ$ ;  $I = -70.3 \pm 2.1^\circ$ .

The Port Waikato samples show very consistent behaviour, with linear decay for both thermal and AF demagnetisation above 180–210 °C and 10–12 mT, respectively (Figure 3E,F). The stable remanent magnetisation component is removed at 480–510 °C or at 60–80 mT. The 48 specimens from Port Waikato yield a mean direction  $D = 346.2 \pm 9.5^\circ$  and  $I = -74.1 \pm 2.9^\circ$ , and a tilt-corrected direction  $D = 61.0 \pm 4.9^\circ$  and  $I = -54.6 \pm 4.0^\circ$  (Figure 4E,F, Table 2). Four samples are rejected by the 45° cut-off and, again, all samples indicate normal polarity.

All sites have an  $A_{95}$  value between  $A_{95_{\min}}$  and  $A_{95_{\max}}$  (Table 2), which suggests that the scatter in the individual datasets can be simply explained by paleosecular variation (Deenen et al. 2011),

### Fold test


We performed a fold test for all three localities, following the eigenvector bootstrap approach of Tauxe and Watson (1994) in [www.paleomagnetism.org](http://www.paleomagnetism.org). The eight samples of AG2\* that form a separate cluster (Figure 4A,B) are not taken into account in the Awakino Gorge fold test in the absence of significant bedding tilt differences. The remaining Awakino Gorge directions clearly fail the fold test, with highest clustering around 0% unfolding, ranging [–8%, +6%] at the 95% level (Figure 6A).

The fold test performed on Kiritehere samples shows a less straightforward result, and highest clustering is found around an unfolding percentage of 44%, ranging [15–73%] (Figure 6B). This may be the result of remagnetisation during folding, but it may also be the result of a new phase of folding after a first post-folding remagnetisation occurred (Tauxe and Watson 1994). Therefore, the data fail the fold test and tectonically corrected directions for Kiritehere sample sites cannot be used for tectonic analysis.

The Port Waikato samples were all taken at one site at an outcrop with a very constant bedding orientation and consequently a fold test cannot be performed on the directions of the Port Waikato locality.

### Discussion


The most striking result from our paleomagnetic analysis is the almost exclusively normal polarity held within samples from all three localities, despite the large stratigraphic age range (c. 209–145 Ma) sampled. During accumulation of Murihiku sediments, many polarity reversals took place (Gradstein et al. 2012). Therefore, the almost complete absence of samples with reversed



**Figure 4.** Equal area projections of characteristic remanent magnetisation (ChRM) directions per site. Closed (open) circles indicate lower (upper) hemisphere projection. Boxes give the mean with D/I as declination/inclination, K is the precision parameter of the VGP distribution, Na is the number of accepted samples accepted by the 45° cut-off.

polarity, with the possible exception of AG2\*, implies widespread remagnetisation. The negative fold test of Awakino Gorge demonstrates that the remanent magnetisation signal was acquired after folding of the succession. The outcome of the fold test performed on Kiritehere data demonstrates that these samples do not carry their original (or at least pre-folding)

magnetisation, but the exact timing of the magnetisation relative to the deformation history is more difficult to assess. Therefore, we conclude that most of the sample host rocks have been remagnetised. As a result, the bulk of our data cannot help constrain the Mesozoic paleolatitude of the Murihiku Terrane. However, as the paleomagnetic directions do not correspond to a recent




**Figure 5.** Examples of demagnetisation diagrams in both geographic (no TC, left column) and tectonic (TC, right column) coordinates for samples from sites AG2\* (yellow) and AG2 (blue). For symbols, refer to caption to Figure 3.

geocentric axial dipole (GAD) direction (Figure 4), we explore their value in identifying a tectonic signal since remagnetisation. This implies that we need an estimate of the age of remagnetisation. The meaning of the small set of directions in AG2\* that show a different paleomagnetic result are also discussed.

### Remagnetisation event

The occurrence of widespread remagnetisation in New Zealand has been demonstrated previously by Haston and Luyendyk (1991), Oliver (1994) and Kodama et al. (2007). The age and origin of the remagnetisation event remains unclear, but is of importance for understanding New Zealand's tectonic history. Some

suggestions of the age and origin have been made, but none has been conclusive. Haston and Luyendyk (1991) undertook a paleomagnetic study of samples from the Waipapa Terrane and compared their computed paleomagnetic pole (69.7°S, 150.7°E) to a synthetic apparent polar wander path of New Zealand. They concluded that the remagnetisation event most likely occurred between 12 Ma and 58 Ma. They prefer a 25 Ma age of remagnetisation, caused by a major change in the tectonic regime. The change in tectonic regime has been associated with: (i) emplacement of ophiolites across northern Northland Peninsula at that time (Ballance and Spörli 1979), (ii) inception of subduction along the Hikurangi margin (e.g. Cole and Lewis 1981; Kamp 1999) and (iii) initiation of



**Figure 6.** Results of the eigenvector bootstrap ( $N_b = 1000$ ) fold test according to Tauxe and Watson (1994) performed on the Awakino (A) and Kiritehere (B) data sets. Shaded area shows the cumulative distribution function of the maximum unfolding at the 95% bootstrap confidence level. The first 25 bootstraps are shown.


arc volcanism (e.g. Mortimer et al. 2010). Remagnetisation in the Waipapa Terrane may be due to fluid migration associated with those events. Oliver (1994) undertook a paleomagnetic study of rocks from the Kawhia and Marokopa areas within the Murihiku Terrane north of Kiritehere. He acquired exclusively normal polarity magnetic directions, in agreement with our findings. He obtained a paleomagnetic pole from the remagnetised directions ( $57.2^\circ\text{S}$ ,  $163.2^\circ\text{E}$ ), which is consistent with the c. 80–90 Ma position predicted by the apparent polar wander path of Australia, as well as a Cretaceous pole determined for Marie Byrd Land (West Antarctica) by Grindley and Oliver (1983). Oliver (1994) thus suggested that remagnetisation took place around 85 Ma, possibly linked to onset of Tasman Sea spreading around that time (Weissel et al. 1977; Weissel and Hayes 1977; Gaina et al. 1998). Kodama et al. (2007) concluded that remagnetisation must have occurred no later than Late Paleocene–Eocene, since they argued for a primary magnetic signal in Late Paleocene–Eocene sedimentary rocks in Kaiwhata Stream.

The timing of folding of the Kawhia syncline and its parasitic smaller-scale folds, has been constrained from thermal history modelling of apatite fission track data obtained from samples collected from the Port Waikato area (Kamp and Liddell 2000). These data indicate

cooling of 110–120 °C along the eastern limb of the syncline by c. 100 Ma, indicating that folding occurred during the early Cretaceous.

Furthermore, the remagnetisation of strata within the Kawhia syncline probably occurred before uplift and erosion of the Kawhia syncline succession, which accompanied or followed deformation of the succession. This can be reasoned from the requirement that remagnetisation is thought to require burial temperatures higher than those in the near subsurface following the bulk of the erosion. The timing of uplift and erosion of Murihiku Terrane is constrained stratigraphically to lie between: (i) the latest Jurassic, which is the age of the Huriwai Group, forming the youngest strata involved in the syncline and hence giving a maximum age when peak burial heating was experienced by the strata; and (ii) the Late Eocene, which is the age of the overlying Te Kuiti Group (Nelson 1978; Edbrooke 2005; Kamp et al. 2014). From this, we conclude that remagnetisation most probably occurred after 100 Ma and before 40 Ma.

We confined the estimated age of remagnetisation further by plotting our obtained geomagnetic directions on the apparent polar wander path of New Zealand. To this end, we calculated the Global Apparent Polar Wander Path of Torsvik et al. (2012) in the coordinates of New Zealand north of the Alpine Fault (northern Zealandia), using the Euler rotations provided in the plate reconstruction of Müller et al. (2016). We used the thereto designed tool at [www.paleomagnetism.org](http://www.paleomagnetism.org) as described in Li et al. (2017). Comparison between the paleolatitudes of reference location ( $38.359^\circ\text{S}$ ,  $174.802^\circ\text{E}$ ) calculated from the directions in geographic coordinates for both Port Waikato and Kiritehere and the predicted Apparent Polar Wander Path for northern Zealandia, suggests an age of remagnetisation of  $80 \pm 8$  Ma (Figure 7, black bar on horizontal axis). Because all our samples carry a normal polarity signal, resetting of the magnetic signal must have taken place during a normal polarity interval. Based on the  $80 \pm 8$  Ma fit, remagnetisation then likely occurred during the Cretaceous Normal Superchron (126–83.6 Ma), or possibly during younger chrons such as 33n (79.9–74.3 Ma) (Gradstein et al. 2012). Because of its shallower inclination ( $I = -58.2^\circ$ ) compared with our other sites, the corresponding paleolatitude of Awakino Gorge differs considerably from those of Port Waikato and Kiritehere, and would be close to the present-day latitude of New Zealand (Figure 7). This could indicate that the remagnetisation is a relatively young (Eocene) to very recent field overprint. However, if the samples of Awakino Gorge were overprinted by a recent field, a declination of  $0^\circ$  would be expected, which is not observed (Figure 4). We consider a possible explanation for this discrepancy below.



**Figure 7.** Diagram of the paleolatitudes (geographic coordinates) of sampling localities AG, KI, PW and the AG locality corrected for a 20° dip towards the east vs. the Apparent Polar Wander Path (APWP) of northern Zealandia. From the uncertainties in paleolatitude we deduce an age of remagnetisation of c.  $83 \pm 5$  Ma. The paleolatitude of Awakino Gorge significantly deviates from the paleolatitudes of Kiritehere and Port Waikato, but becomes very similar after the E-dip correction.

We note that an c. 80 Ma age of remagnetisation coincides with the early stages of seafloor spreading in the Tasman Sea (Gaina et al. 1998), as previously noted by Oliver (1994). Additionally, remagnetisation may be related to the end of subduction along the Gondwana margin of New Zealand. The timing of the end of subduction is subject to debate and is generally thought to have occurred around 105–100 Ma (Bradshaw 1989; Luyendyk 1995; Davy et al. 2008; Matthews et al. 2012), but ages around 85 Ma have been proposed (Mazengarb and Harris 1994; Kamp and Liddell 2000; Vry et al. 2004), for example based on zircon fission track analysis of the youngest rocks in the toe of the Torlesse Complex (Kamp 1999, 2000).

### Post-remagnetisation tectonics affecting the Awakino Gorge area?

Given the evidence above for pervasive and widespread remagnetisation in the Kawhia syncline, as well as by previous studies in the North Island and Marlborough, South Island (Haston and Luyendyk 1991; Oliver 1994; Kodama et al. 2007), it is remarkable that *in situ* directions for samples from Awakino Gorge differ significantly from those obtained from the Port Waikato and Kiritehere sections. As we have already identified, if the *in situ* direction in AG samples is an overprint direction, it must have been acquired sometime between the Eocene and present day (Figure 7). This is very unlikely, however, given the large difference of  $> 100^\circ$  between the GAD declination and the declination observed at Awakino Gorge (Figure 4). We therefore seek an alternative explanation for the lower inclination and the declination of Awakino Gorge data.

An inferred Late Cretaceous overprint would yield an observed inclination of around  $-74^\circ$  at the reference


location ( $38.359^\circ\text{S}$ ,  $174.802^\circ\text{E}$ ), identical within error to the values reported here for the Port Waikato and Kiritehere sample sites (Table 2). Because the average observed inclination for the Awakino Gorge samples is  $-58.2 \pm 3.6^\circ$ , the sample host strata have been subject to inclination lowering of c.  $15^\circ$ .

The western limb of the syncline in the Awakino Gorge area has been accentuated by Late Oligocene and Early Miocene reverse displacement on the Manganui Fault (Kamp et al. 2004). Because this fault lies to the west of the Awakino Gorge paleomagnetic sampling sites, it probably increased the eastward dip of the beds we sampled by 20° as they lie in the hanging wall of the fault. This post-remagnetisation deformation may have contributed to the inclination shallowing of the magnetisation in these rocks compared with those at the Kiritehere and Port Waikato sample sites.

Because the 20° eastward tilting occurred after remagnetisation, we have to apply a 000/20 bedding tilt correction to our data from Awakino Gorge. This leads to  $I = -77 \pm 2.9^\circ$  and  $D = 121.0 \pm 12.0^\circ$  (for full parameters see Table 2). The newly obtained inclination is far more consistent with the data of Kiritehere and Port Waikato. We can now also use the Awakino Gorge data to obtain the age of remagnetisation, which is now constrained to  $83 \pm 5$  Ma (Figure 7, yellow bar on horizontal axis).

### Rotation of Murihiku Terrane

Another interesting feature is that the declination of the Kiritehere data set corresponds well to the 80 Ma declination of New Zealand, whereas the declinations of both Port Waikato and Awakino Gorge deviate (Figure 8). This suggests that parts of Murihiku Terrane have been rotated since the widespread remagnetisation. Because our Kiritehere data record a



**Figure 8.** Diagram of declinations (geographic coordinates) of our sampling sites and the Apparent Polar Wander Path (APWP) of northern Zealandia. The declination of Kiritehere fits well with our inferred age of remagnetisation of  $83 \pm 5$  Ma, but declinations of Port Waikato and Awakino Gorge deviate, indicating respectively  $39.4 \pm 9.5^\circ$  counterclockwise and  $77.3 \pm 5.0^\circ$  (uncorrected for  $20^\circ$  eastward dip) and  $95.8 \pm 12.0^\circ$  (corrected for  $20^\circ$  eastward dip) clockwise rotations for these areas.


declination very similar to the declination of New Zealand at 80 Ma, it suggests that this part of the Murihiku Terrane has not been rotated relative to New Zealand since remagnetisation. Other paleomagnetic studies of Murihiku Terrane strata in the Kawhia area north of Kiritehere also suggested that the Murihiku Terrane did not experience significant amounts of vertical axis rotation since remagnetisation (Oliver 1994; Kodama et al. 2007) (Figure 9).

By contrast, our results suggest that the basement in Awakino Gorge rotated about  $77.3 \pm 5.0^\circ$  clockwise, whereas the basement in the Port Waikato area has rotated c.  $39.4 \pm 9.5^\circ$  counterclockwise since remagnetisation (Figure 9). After correcting the site data from Awakino Gorge for the  $20^\circ$  eastward dip, the rotation of the basement succession in Awakino Gorge increases to c.  $95.8 \pm 12.0^\circ$ .

Previous paleomagnetic studies do indicate large-scale Neogene clockwise rotation along the Hikurangi margin, related to deformation considered to be associated with active subduction to the east of New Zealand (Wright and Walcott 1986; Rowan et al. 2005; Lamb 2011). These rotations may have affected the southern part of the Murihiku Terrane (i.e. Awakino Gorge) as well. However, previous paleomagnetic studies carried out on Oligocene calcareous siltstone (Whaingaroa Formation) exposed at the entrance to Bexley Station in Awakino Gorge (Lamb 2011, figure 4), record clockwise rotations up to  $30^\circ$  that had ceased by the Early Miocene (Mumme and Walcott 1985, cited in Lamb 2011). No paleomagnetic evidence exists for later rotations in this area.

We propose that rotation of the Murihiku Terrane in the Kawhia syncline is not directly related to modern Australia–Pacific plate boundary zone deformation. Clockwise rotation in the Awakino Gorge area is thought to have ceased before the c. 30–25 Ma inception of the Alpine Fault (Kamp 1986; Lamb 2011),

and it would also fail to explain the counterclockwise rotation at Port Waikato. We suggest that the observed rotation may have been related to Cretaceous bending of basement terranes expressed in the New Zealand orocline, the origin of which remains unknown. The timing of bending is also subject to debate. Essentially, there are three scenarios regarding the age of the New



**Figure 9.** Map showing declinations of our sites and those of Oliver (1994) and Kodama et al. (2007) with respective confidence parachutes ( $\Delta D_x$ ). Declinations of Oliver (1994) and Kodama et al. (2007) have been acquired by recalculating their published data using [www.paleomagnetism.org](http://www.paleomagnetism.org)

Zealand orocline (Bradshaw et al. 1996, see also Lamb et al. 2016): (i) oroclinal bending is older than Late Cretaceous; (ii) large-scale bending occurred exclusively during the Cenozoic; and (iii) the basement terranes of New Zealand were partially bent before the Cenozoic, but a substantial amount of rotation occurred during the Cenozoic.

Our results suggest that considerable rotation occurred since the Late Cretaceous ( $83 \pm 5$  Ma), which would argue against the first option given above. Because we cannot rule out the possibility of partially bent terranes before c. 83 Ma, suggestions (ii) and (iii) are both considered possible. There is no paleomagnetic evidence for any rotations that occurred in this area since the Miocene, therefore the bulk of bending occurred between  $83 \pm 5$  Ma and c. 20 Ma.

The rotation of Port Waikato with respect to Kiriterehere is fairly consistent with the overall strike of the Kawhia syncline. The difference in the general strike of the Kawhia syncline at Kiriterere (c.  $10^\circ$ ) and Port Waikato (c.  $335^\circ$ ) is c.  $35^\circ$  (see Figure 1, based on eastern limb), which corresponds well to our inferred rotation (c.  $40^\circ$ ). However, the large rotation of the basement succession in Awakino Gorge with respect to Kiriterehere cannot be explained in the same way. Therefore, a substantial part of the rotation at Awakino Gorge must be a local phenomenon. At this point we cannot explain the origin of the rotation at Awakino Gorge, but our data clearly indicate its occurrence.

Additional paleomagnetic studies can usefully be carried out to place tighter constraints on the origin and timing of rotation of the Murihiku Terrane in the Kawhia syncline, and the possible relation to bending of basement terranes. Comparing the amount of rotation predicted from the declination of Cenozoic rocks from the Port Waikato and Awakino Gorge areas with the declinations of this study allows estimation of the timing of rotation. When the age of rotation is more tightly constrained, a hypothesis can be formulated about the origin of rotation in sample host rocks in both the Awakino Gorge and Port Waikato areas, and the possible relation to bending of the basement terranes can be investigated more completely. Importantly, our study shows that remagnetised Murihiku Terrane sedimentary rocks widely exposed in the central-western North Island, are excellent recorders of a Cretaceous paleomagnetic field yet to be completely defined, but which can nevertheless be used to deduce post-Late Cretaceous regional vertical axis rotation and horizontal-axis tilt histories.

### **A primary, Jurassic magnetisation at Awakino Gorge?**

As mentioned previously, eight samples collected from site AG2 (Early Jurassic, Toarcian age, 181–176 Ma) in

Awakino Gorge record a magnetic component that is distinct from all other samples. This is a high-temperature component that provides an *in situ* inclination of only  $5.8^\circ$ . If acquired after bedding tilt, as with the regional ChRM interpreted from all other samples, this would require a latitude of magnetisation close to the equator. However, New Zealand has not been close to the equator since the Toarcian. Rather, it was located very close to the South Pole. It is therefore highly unlikely that these samples acquired their magnetisation after tilting.


We therefore tentatively assume that these samples have retained their pre-tilt magnetisation and escaped complete remagnetisation. We thus correct the AG2\* directions for bedding tilt to obtain the original inclination. This gives a direction  $D = 316 \pm 27.0^\circ$  and  $I = 74.9 \pm 7.5^\circ$  (Figure 4, Table 2). This would imply that Murihiku Basin was located at a paleolatitude of  $61.6^\circ\text{N}$  during the Toarcian, which is not possible. Therefore, we assume that these rocks record a reversed magnetic signal and that the Murihiku forearc basin was located at  $61.6 \pm 12^\circ\text{S}$  during the Toarcian.

Because we previously inferred that Awakino Gorge has been subject to a Late Oligocene to Early Miocene  $20^\circ$  tilt, we have to apply an additional bedding tilt correction. The corrected geomagnetic directions are  $D = 308.3 \pm 27.6^\circ$  and  $I = 75.2 \pm 7.5^\circ$ , corresponding to a paleolatitude of  $62.1 \pm 12^\circ\text{S}$  (Table 2).

This paleolatitude is in agreement with that of Grindley et al. (1980), which is the only other paleomagnetic study that may have found a paleolatitude from a primary magnetic signal of Murihiku Terrane rocks. They obtained a paleolatitude of  $66^\circ\text{S}$  for the Glenham Porphyry of Early Jurassic (c. 190 Ma) age in the Southland syncline part of the Murihiku Terrane. Both studies therefore imply that the Murihiku Terrane was located in the vicinity of the New Zealand sector of the eastern Gondwana margin during sediment accumulation.

When compared with the apparent polar wander path of northern Zealandia, our calculated paleolatitude is located within c.  $20^\circ$  expected for Zealandia in the Toarcian (Figure 10). This would thus provide a first-order estimate that the intra-oceanic arc to which the Murihiku forearc basin was paired, was probably located within 1000–2000 km, but possibly up to 3500 km, north of the contemporary Gondwana margin (Figure 11).

The similarity in paleolatitude from our small dataset and the one of Grindley et al. (1980) is promising, but the results of both studies are subject to uncertainties. The findings of Grindley et al. (1980) have been called doubtful due to uncertainty in tilt-corrections for volcanic rocks (Kodama et al. 2007), while our data set is very small and lacks independent field tests. Therefore, we emphasise the tentative nature of our interpretation, and a larger data set is needed to



**Figure 10.** Apparent Polar Wander Path (APWP) of northern Zealandia and the paleolatitude of site AG2\*, first corrected for the eastward dip of 20° (plunge correction), and subsequently tilt corrected with the plunge corrected bedding plane.

put better constraints on the original paleolatitude of the Murihiku forearc basin.

Finding the original signal has proven to be extremely difficult due to the widespread magnetic overprint. Nevertheless, our data show that such an effort is not futile and provides a lead to identify primary magnetisations in Murihiku rocks. It would be worthwhile to carry out extensive sampling of the rock succession exposed in Awakino Gorge and in Glenham Porphyry to increase the data set and quantitatively constrain the Mesozoic plate kinematic history of the Gondwana margin and adjacent intra-oceanic arcs of New Zealand.


## Conclusions

To aid kinematic restoration of the Mesozoic and Cenozoic plate tectonic and orogenic history of part of New Zealand, we performed a paleomagnetic study on rocks from the Upper Triassic to Lower Jurassic of the Murihiku forearc basin in the central–western North Island, New Zealand. Our results confirm previous interpretations of widespread pervasive remagnetisation throughout the North Island in the form of exclusively normal polarities – despite accumulation of the rocks during times of high reversal frequency – and negative fold tests. We show demagnetisation results for 248 samples from three localities at Awakino Gorge, Port Waikato, and Kiritehere Beach. The latter two record similar *in situ* inclinations of c.  $-74^\circ$ , corresponding to a latitude of  $61^\circ\text{S}$  during post-tilt remagnetisation. Based on comparison of the expected paleolatitude of these samples with the apparent polar wander path of New Zealand, we estimate remagnetisation to have occurred around  $83 \pm 5$  Ma.

We show that remagnetised directions in Murihiku Terrane sediments provide an excellent datum for post-remagnetisation studies of regional tilt and vertical axis rotation. We show that the lower inclinations in Awakino Gorge paleomagnetic data compared with data for the more northern sample localities are consistent with a regional eastward tilt of c.  $20^\circ$  as a result of reverse displacement on Manganui Fault.

The Murihiku Terrane has, in addition, been subjected to significant rotation since remagnetisation. Port Waikato has been rotated  $39.4 \pm 9.5^\circ$  counterclockwise and Awakino Gorge has been rotated  $77.3 \pm 5.0^\circ$  clockwise ( $95.8 \pm 12.0^\circ$  when corrected for eastward tilt), whereas Kiritehere rocks have remained stable with respect to Zealandia. We suggest that these rotations can be used to reconstruct the tectonic bending of basement terranes, which must have occurred, at least in part, after the remagnetisation event estimated at  $83 \pm 5$  Ma.

Only 8 of the 93 demagnetised Awakino Gorge samples, all at the Awakino Gorge 2 site, retain a



**Figure 11.** Map showing the possible locations (yellow disc, latitudes 50°S–75°S) of the Murihiku forearc basin based on site AG2\* (reference location 38.373°S, 174.711°E). The grey area is the position of Gondwana based on the reconstruction of Müller et al. (2016) in the paleomagnetic reference frame of Torsvik et al. (2012). Present-day coastlines are for reference only. The blue zone indicates the uncertainty of the position of northern Zealandia within Gondwana. The overlap between the paleolatitude belt and the blue zone (green belt) indicates the possible locations of Murihiku forearc basin if it was part of the Gondwana margin.

high-temperature, high-coercivity magnetic component that may represent a primary magnetisation. *In situ* inclinations of only 5° would require equatorial latitudes of magnetisation, which is not possible considering plate reconstruction constraints. A primary magnetisation cannot be confirmed by independent field tests, but pre-bedding tilt directions of these eight samples suggest a paleolatitude for Murihiku Terrane of  $62.1 \pm 12^\circ\text{S}$  during the Early Jurassic (Toarcian), c. 15° north of the New Zealand sector of the contemporary eastern Gondwana Margin.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

LMB acknowledges NWO grant 824.01.004. PJKK acknowledges New Zealand MBIE UOWX1501 funding (contract number CONT- 42907-EMTR-UOW). DJJvH acknowledges ERC Starting Grant 306810 (SINK) and NWO Vidi grant 864.11.004.

## References

- Aita Y, Spörli KB. 1992. Tectonic and paleobiogeographic significance of radiolarian microfaunas in the Permian to Mesozoic basement rocks of the North Island, New Zealand. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 96:103–125.
- Ballance PF, Campbell JD. 1993. A Murihiku arc-related basin of New Zealand (Triassic–Jurassic). In: Ballance PF, editor. *South Pacific sedimentary basins*. Amsterdam: Elsevier; p. 21–33.
- Ballance PF, Heming RF, Sameshima T. 1980. Petrography of the youngest known Murihiku Supergroup, New Zealand: latest Jurassic arc volcanism on the southern margin of Gondwana. In: Cresswell MM, Vella P, editors. *Gondwana five*. Rotterdam: Balkema; p. 161–165.
- Ballance PF, Spörli KB. 1979. Northland allochthon. *Journal of the Royal Society of New Zealand*. 9:259–275.
- Bradshaw JD. 1989. Cretaceous geotectonic patterns in the New Zealand region. *Tectonics*. 8:803–820. doi:10.1029/TC008i004p00803.
- Bradshaw JD. 1994. Brook street and Murihiku terranes of New Zealand in the context of a mobile south Pacific Gondwana margin. *Journal of South American Earth Sciences*. 7:325–332.
- Bradshaw JD, Weaver SD, Muir RJ. 1996. Mid-Cretaceous oroclinal bending of New Zealand terranes. *New Zealand Journal of Geology and Geophysics*. 39:461–468.
- Briggs RM, Middleton MP, Nelson CS. 2004. Provenance history of a late Triassic–Jurassic Gondwana margin fore-arc basin, Murihiku terrane, north island, New Zealand: petrographic and geochemical constraints. *New Zealand Journal of Geology and Geophysics*. 47:589–602.
- Butler RF. 1992. *Paleomagnetism: magnetic domains to geologic terrains*. Boston (MA): Backwell Scientific Publications.
- Campbell HJ, Mortimer N, Turnbull IM. 2003. Murihiku supergroup, New Zealand: redefined. *Journal of the Royal Society of New Zealand*. 33:85–95. doi:10.1080/03014223.2003.9517722.
- Challinor AB. 2001. Stratigraphy of Tithonian (Ohauan-Puaruan) marine beds near Port Waikato, New Zealand, and a redescription of *Belemnopsis aucklandica* (Hochstetter). *New Zealand Journal of Geology and Geophysics*. 44:219–242. doi:10.1080/00288306.2001.9514936.
- Cole JW, Lewis KB. 1981. Evolution of the Taupo–Hikurangi subduction system. *Tectonophysics*. 72:1–21.
- Coombs DS. 1954. The nature and alteration of some Triassic sediments from Southland, New Zealand. *Transactions of the Royal Society of New Zealand*. 82:65–109.
- Coombs DS, Cook NDJ, Campbell JD. 1992. The park volcanics group: field relations of an igneous suite emplaced in the Triassic–Jurassic Murihiku Terrane, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics*. 35:337–351. doi:10.1080/00288306.1992.9514527.
- Dankers PHM. 1978. *Magnetic properties of dispersed natural iron-oxides of known grain-size* [PhD Thesis]. University of Utrecht; p. 142.
- Davy B, Hoernle K, Werner R. 2008. Hikurangi plateau: crustal structure, rifted formation, and Gondwana subduction history. *Geochemistry, Geophysics, Geosystems*. 9:Q07004. doi:10.1029/2007GC001855.
- Deenen MHL, Langereis CG, van Hinsbergen DJJ, Biggin AJ. 2011. Geomagnetic secular variation and the statistics of palaeomagnetic directions. *Geophysical Journal International*. 186:509–520.
- Deenen MHL, Langereis CG, Van Hinsbergen DJJ, Biggin AJ. 2014. Erratum: geomagnetic secular variation and the statistics of palaeomagnetic directions. *Geophysical Journal International*. 197:643.
- Edbrooke SW. (compiler). 2001. *Geology of the Auckland area: scale 1:250,000*. Lower Hutt: Institute of Geological & Nuclear Sciences Limited. Institute of Geological & Nuclear Sciences 1:250,000 geological map 3. 74 p. + 1 folded map.
- Edbrooke SW. (compiler). 2005. *Geology of the Waikato area: scale 1:250,000*. Lower Hutt: Institute of Geological & Nuclear Sciences Limited. Institute of Geological & Nuclear Sciences 1:250,000 geological map 3. 74 p. + 1 folded map.
- Edbrooke SW, Heron DW, Forsyth PJ, Jongens R. 2015. *Geology map of New Zealand 1: 1 000 000*. GNS Science Map, 2.
- Fisher R. 1953. Dispersion on a sphere. In: *Proceedings of the Royal Society of London A: mathematical, physical and engineering sciences*. The Royal Society; p. 295–305.
- Gaina C, Müller RD, Royer JY, Stock J, Hardebeck J, Symonds P. 1998. The tectonic history of the Tasman Sea: a puzzle with 13 pieces. *Journal of Geophysical Research: Solid Earth*. 103:12413–12433.
- Gradstein FM, Ogg JG, Schmitz M, Ogg G, editors. 2012. *The geologic time scale 2012*. Boston (MA): Elsevier.
- Grindley GW, Oliver PJ. 1983. Paleomagnetism of Cretaceous volcanic rocks from Marie Byrd Land, Antarctica. In: Oliver RL, James PR, Jago JB, editors. *Antarctic earth science, Proceedings of the 4th International Symposium on Antarctic Earth Sciences*. Canberra: Australian Academy of Science; p. 573–578.
- Grindley GW, Oliver PJ, Sukroo JC. 1980. Lower Mesozoic position of southern New Zealand determined from paleomagnetism of the Glenham Porphyry, Murihiku Terrane, Eastern Southland. In: Cresswell and Vella, editor. *Gondwana five*. Rotterdam: Balkema; p. 319–326.

- Haston RB, Luyendyk BP. 1991. Paleomagnetic results from the Waipapa Terrane, Northland Peninsula, North Island, New Zealand: tectonic implications from widespread remagnetizations. *Tectonics*. 10:986–994.
- Hay RF. 1967. Sheet 7, Taranaki, 'Geological map of New Zealand 1:250 000'. DSIR Wellington.
- Howell D. 1980. Mesozoic accretion of exotic terranes along the New Zealand segment of gondwanaland. *Geology*. 8:487–491.
- Johnson CL, Constable CG, Tauxe L, Barendregt R, Brown LL, Coe RS, Layer P, Mejia V, Opdyke ND, Singer BS, et al. 2008. Recent investigations of the 0–5 Ma geomagnetic field recorded by lava flows. *Geochemistry, Geophysics, Geosystems*. 9:Q04032. doi:10.1029/2007GC001696.
- Kamp PJJ. 1986. The mid-Cenozoic challenger rift system of western New Zealand and its implications for the age of alpine fault inception. *Geological Society of America Bulletin*. 97:255–281.
- Kamp PJJ. 1987. Age and origin of the New Zealand orocline in relation to alpine fault movement. *Journal of the Geological Society*. 144:641–652.
- Kamp PJJ. 1999. Tracking crustal processes by FT thermochronology in a forearc high (Hikurangi margin, New Zealand) involving Cretaceous subduction termination and mid-Cenozoic subduction initiation. *Tectonophysics*. 307:313–343.
- Kamp PJJ. 2000. Thermochronology of the Torlesse accretionary complex, Wellington region, New Zealand. *Journal of Geophysical Research: Solid Earth*. 105:19253–19272.
- Kamp PJJ, Liddell IJ. 2000. Thermochronology of northern Murihiku Terrane, New Zealand, derived from apatite FT analysis. *Journal of the Geological Society*. 157:345–354.
- Kamp PJJ, Tripathi ARP, Nelson CS. 2014. Paleogeography of late Eocene to earliest Miocene Te Kuiti group, central-western North Island, New Zealand. *New Zealand Journal of Geology and Geophysics*. 57:128–148. doi:10.1080/00288306.2014.904384.
- Kamp PJJ, Vonk AJ, Nelson CS, Hansen RJ, Tripathi A, Hood S, Ngatai M, Hendy AJW. 2004. Constraints on the evolution of Taranaki fault from thermochronology and basin analysis: implications for the Taranaki fault play. 2004 New Zealand Petroleum Conference Proceedings; 7–10th March 2004; Auckland. Crown Minerals. Ministry of Economic Development, Wellington. 25pp. [http://www.med.govt.nz/pubs/publications-03.html#P232\\_7656](http://www.med.govt.nz/pubs/publications-03.html#P232_7656)
- Kear D. 1960. Sheet 4, Hamilton, 'Geological map of NZ 1:250 000'. DSIR Wellington.
- Kirschvink JL. 1980. The least-squares line and plane and the analysis of palaeomagnetic data. *Geophysical Journal International*. 62:699–718.
- Kodama K, Fukuoka M, Aita Y, Sakai T, Hori RS, Takemura A, Campbell HJ, Hollis CJ, Grant-Mackie JA, Spörli KB. 2007. Paleomagnetic results from arrow rocks in the framework of paleomagnetism in pre-Neogene rocks from New Zealand. In: Spörli B, Takemura A, Hori RS, editors. The oceanic Permian/Triassic boundary sequence at arrow rocks (Oruatemanu), Northland, New Zealand. Wellington: GNS Science Monograph; p. 177–196.
- Koymans MR, Langereis CG, Pastor-Galan D, van Hinsbergen DJJ. 2016. Paleomagnetism.org: An online multi-platform open source environment for paleomagnetic data analysis. *Computers and Geosciences*. 93:127–137.
- Lamb S. 2011. Cenozoic tectonic evolution of the New Zealand plate-boundary zone: A paleomagnetic perspective. *Tectonophysics*. 509:135–164.
- Lamb S, Mortimer N, Smith E, Turner G. 2016. Focusing of relative plate motion at a continental transform fault: Cenozoic dextral displacement >700 km on New Zealand's alpine fault, reversing >225 km of late Cretaceous sinistral motion. *Geochemistry, Geophysics, Geosystems*. 17:1197–1213. doi:10.1002/2015GC006225.
- Li S, Advokaat EL, van Hinsbergen DJJ, Koymans MR, Deng C, Zhu R. 2017. Paleomagnetic constraints on the Mesozoic-Cenozoic paleolatitudinal and rotational history of Indochina and South China: review and updated kinematic reconstruction. *Earth-Science Reviews*. 171:58–77.
- Luyendyk BP. 1995. Hypothesis for Cretaceous rifting of east Gondwana caused by subducted slab capture. *Geology*. 23:373–376.
- MacKinnon TC. 1983. Origin of the Torlesse terrane and coeval rocks, South Island, New Zealand. *Geological Society of America Bulletin*. 94:967–985.
- Matthews KJ, Seton M, Müller RD. 2012. A global-scale plate reorganization event at 105–100Ma. *Earth and Planetary Science Letters*. 355–356:283–298. doi:10.1016/j.epsl.2012.08.023.
- Mazengarb C, Harris DHM. 1994. Cretaceous stratigraphic and structural relations of Raukumara Peninsula, New Zealand: stratigraphic patterns associated with the migration of a thrust system. *Annales Tectonicae*. 8:100–118.
- McFadden PL, McElhinny MW. 1988. The combined analysis of remagnetization circles and direct observations in palaeomagnetism. *Earth and Planetary Science Letters*. 87:161–172.
- Mortimer N. 2004. New Zealand's geological foundations. *Gondwana Research*. 7:261–272.
- Mortimer N. 2014. The oroclinal bend in the South Island, New Zealand. *Journal of Structural Geology*. 64:32–38. doi:10.1016/j.jsg.2013.08.011.
- Mortimer N, Gans PB, Palin JM, Meffre S, Herzer RH, Skinner DNB. 2010. Location and migration of Miocene–Quaternary volcanic arcs in the SW Pacific region. *Journal of Volcanology and Geothermal Research*. 190:1–10. doi:10.1016/j.jvolgeores.2009.02.017.
- Mortimer N, Rattenbury MS, King PR, Bland KJ, Barrell DJA, Bache F, Begg JG, Campbell HJ, Cox SC, Crampton JS, et al. 2014. High-level stratigraphic scheme for New Zealand rocks. *New Zealand Journal of Geology and Geophysics*. 57:402–419. doi:10.1080/00288306.2014.946062.
- Mullender TAT, Van Velzen AJ, Dekkers MJ. 1993. Continuous drift correction and separate identification of ferrimagnetic and paramagnetic contributions in thermomagnetic runs. *Geophysical Journal International*. 114:663–672.
- Müller RD, Seton MS, Zahirovic S, Williams SE, Matthews KJ, Wright NM, Shephard GE, Maloney KT, Barnett-Moore N, Hosseinpour M, et al. 2016. Ocean basin evolution and global-scale plate reorganization events since Pangea breakup. *Annual Review of Earth and Planetary Sciences*. 44:107–138. doi:10.1146/annurev-earth-060115-012211.
- Mumme TC, Walcott RI. 1985. Paleomagnetic studies at geophysics division, 1980–1983. Geophysics Division, Department of Scientific and Industrial Research 204.
- Nelson CS. 1978. Stratigraphy and paleontology of the Oligocene Te Kuiti group, Waitomo county, South

- Auckland, New Zealand. *New Zealand Journal of Geology and Geophysics*. 21:553–594. doi:10.1080/00288306.1978.10424086.
- Oliver PJ. 1994. The tectonic significance of paleomagnetic results from the Triassic and Jurassic Murihiku sedimentary rocks of the Kawhia region, North Island, New Zealand. In: Van der Lingen GJ, Swanson KM, Muir RJ, editors. *Evolution of the Tasman Sea basin: proceedings of the Tasman Sea conference*. Rotterdam: Balkema; p. 67–82.
- Passier HF, de Lange GJ, Dekkers MJ. 2001. Rock-magnetic properties and geochemistry of the active oxidation front and the youngest sapropel in the Mediterranean. *Geophysical Journal International*. 145:604–614.
- Retallack GJ. 1987. Triassic vegetation and geography of the New Zealand portion of the Gondwana supercontinent. In: McKenzie ED, editor. *Gondwana six: stratigraphy, sedimentology and paleontology*. Washington (DC): American Geophysical Union; p. 29–39.
- Roser BP, Coombs DS, Korsch RJ, Campbell JD. 2002. Whole-rock geochemical variations and evolution of the arc-derived Murihiku Terrane, New Zealand. *Geological Magazine*. 139:665–685. doi:10.1017/S0016756802006945.
- Rowan CJ, Roberts AP, Rait GJ. 2005. Relocation of the tectonic boundary between the Raukumara and Wairoa Domains (East Coast, North Island, New Zealand): implications for the rotation history of the Hikurangi margin. *New Zealand Journal of Geology and Geophysics*. 48:185–196.
- Spörli KB. 1987. Development of the New Zealand microcontinent. In: Monger JWH, Francheteau J, editors. *Circum-pacific orogenic belts and evolution of the Pacific Ocean basin (geodynamimc series 18)*. Washington (DC): American Geophysical Union; p. 115–132.
- Sutherland R. 1999. Cenozoic bending of New Zealand basement terranes and alpine fault displacement: a brief review. *New Zealand Journal of Geology and Geophysics*. 42:295–301. doi:10.1080/00288306.1999.9514846.
- Tauxe L, with contributions from Butler RF, Van der Voo R, Banerjee SK. 2010. *Essentials of paleomagnetism*. Berkeley: University of California Press.
- Tauxe L, Watson GS. 1994. The fold test: an Eigen analysis approach. *Earth and Planetary Science Letters*. 122:331–341.
- Torsvik TH, van der Voo R, Preeden U, Mac Niocail C, Steinberger B, Doubrovine PV, van Hinsbergen DJJ, Domeier M, Gaina C, Tohver E, et al. 2012. Phanerozoic polar wander, palaeogeography and dynamics. *Earth-Science Reviews*. 114:325–368.
- Vry JK, Baker J, Maas R, Little TA, Grapes R, Dixon M. 2004. Zoned (Cretaceous and Cenozoic) garnet and the timing of high grade metamorphism, Southern Alps, New Zealand. *Journal of Metamorphic Geology*. 22:137–157. doi:10.1111/j.1525-1314.2004.00504.x.
- Weissel JK, Hayes DE. 1977. Evolution of the Tasman Sea reappraised. *Earth and Planetary Science Letters*. 36:77–84.
- Weissel JK, Hayes DE, Herron EM. 1977. Plate tectonics synthesis: the displacements between Australia, New Zealand, and Antarctica since the late Cretaceous. *Marine Geology*. 25:231–277.
- Wright IC, Walcott RI. 1986. Large tectonic rotation of part of New Zealand in the last 5 Ma. *Earth and Planetary Science Letters*. 80:348–352.
- Zijderveld JDA. 1967. A.C. demagnetization of rocks: analysis of results. In: Collinson DW, Creer KM, Runcorn SK, editors. *Methods in paleomagnetism*. Amsterdam: Elsevier; p. 254–286.